PROTECT SENSATION® UK-2016-1048

A ready-to-use, 2 in 1 combination of fresh bait and wax block bait containing 0.05 g/kg bromadiolone

Rodenticide for professional use indoors and outdoors (around buildings), including public areas, against the brown rat (*Rattus norvegicus*) and the house mouse (*Mus musculus*)

The rodenticide wax block bait contains denatonium benzoate, an extremely bitter substance.

PROTECT SENSATION®

Ready to use bait containing 0.05 g/kg bromadiolone

Danger

May damage the unborn child

Causes damage to the blood system through prolonged or repeated exposure

Keep out of reach of children.

Obtain special instructions before use.

Do not handle until all safety precautions have been read and understood

Do not get in eyes, on skin, or on clothing.

Do not eat, drink or smoke when using this product.

Wear protective gloves/protective clothing/eye protection/face protection

IF SWALLOWED: Immediately call a POISON CENTRE or doctor

IF exposed or concerned: Get medical advice/attention.

Store locked up.

Dispose of contents and container in accordance with national regulations.

To avoid risks to human health and the environment, comply with the instructions for use

PRECAUTIONS:

To be used only by professional users holding certification demonstrating compliance with UK rodenticide stewardship regime requirements. Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence.

Refer to the CRRU UK Code of Best Practice (or equivalent) for guidance.

When this product is supplied to a user for the control of rodents, it shall only be supplied to a professional user holding certification demonstrating compliance with UK rodenticide stewardship regime requirements. Wash hands and exposed skin before meals and after use. Keep locked up and out of the reach of children.

Unless under the supervision of a pest control operator or other competent person, do not use anticoagulant rodenticides as permanent baits. In most cases, anticoagulant bait should have achieved control within 35 days. Should activity continue beyond this time, the likely cause should be determined.

Prevent access to bait by children, birds and non-target animals (particularly dogs, cats, pigs and poultry). For use only in areas that are inaccessible to infants, children, companion animals and non-target animals.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients. Baits must be securely deposited in a way so as to minimize the risk of consumption by other animals or children. Where possible, secure baits so that they cannot be dragged away.

When the product is being used in public areas, the areas treated must be marked during the treatment period and a notice explaining the risk of primary or secondary poisoning by the anticoagulant, as well as indicating the first measures to be taken in case of poisoning, must be made available alongside the baits. When tamper resistant bait stations are used, they should be clearly marked to show that they contain rodenticides and that they should not be disturbed.

DIRECTIONS FOR USE

PROTECT SENSATION® offers multiple advantages for the user:

- The fresh bait has excellent attractiveness. Rodents can access the fresh bait only if they gnaw off the wax layer of the block, which itself may lead to consumption of a lethal dose.
- The wax outer layer improves the weatherproof features of the bait.
- The bait blocks can be easily and securely fixed in the special bait box, helping to prevent scattering of the bait and minimise exposure to the environment.
- Consumption is easy to monitor and the quantity consumed simple to measure, from the obvious gnawing marks.

PROTECT SENSATION® 2 IN 1 RODENTICIDE BAIT

25 G

Contents: 8 KG

For 24 h emergency information ONLY: call +36 70 637 5436

Bábolna Bio

Place baits manually in the rodent infested area. The bait product can potentially be used under many different circumstances and can be deployed using various means. The method of deployment is dependant on the particular circumstances. The priority is always to exclude non-target exposure as much as possible.

Methods of deployment are bait stations (tamper proof baiting boxes), bait points (a makeshift arrangement using materials and/or the local environment to restrict access to the bait), and loose but inaccessible (using the local environment only to restrict access to the bait).

These methods, in themselves, represent a scale of potential access. The vulnerability (of access by non-target organisms) of a particular site must be assessed in the decision for the deployment method to be used.

Bait can be placed in bait boxes, which may be fixed to the ground. Such bait should be secured in place on the built-in spikes, to minimise removal and dispersal by rodents. Bait may also be placed on trays under a tile, or located in such a way that access by non-target organisms is restricted. The product must never be placed indiscriminately.

The resistance status of the target population should be taken into account when considering the choice of rodenticide to be used. In those areas where evidence of resistance to specific active ingredients is suspected, avoid their use. To control the spreading of resistance, it is advisable to alternate baits containing different anticoagulant active ingredients.

Rate of use: For rat infestations use bait points of 200 g. Place bait points 10 metres apart for low infestations and 7 metres apart for high infestations.

For mouse infestations use one bait point of 40 - 50 g per 5 m^2.

Death of the rodents starts after 3-5 days from first consumption. Make regular inspections of the bait points (every 7 – 10 days) and replace any bait that has been eaten by rodents, damaged by water or contaminated by dirt. If bait is totally consumed then the quantity of the bait at each bait point should be replenished.

First aid: Bromadiolone is an anticoagulant which may produce bleeding; this may be delayed for several days after exposure. If there is no active bleeding the INR (prothrombin time) should be measured on presentation and 48-72 hours after exposure. If the INR is greater than 4, administer Vitamin K1 (phytomenadione) 5-10 mg by slow intravenous injection (100 μg/kg body weight for a child). Treatment with phytomenadione (orally or intravenously) may be required for several weeks. The advice of the National Poisons Information Service (http://www.npis.org/) should be sought, particularly if active bleeding occurs. In case of accident, or if you feel unwell, seek medical advice immediately (show the label where possible). In case of contact with skin, wash with soap and water. Remove and launder any contaminated clothing.

In case of contact with eyes, remove contact lenses if present and rinse the eye slowly and gently with water for 15-20 minutes. Seek medical advice immediately.

In case of ingestion or if swallowed seek medical advice immediately.

Antidote vitamin K1 (under medical supervision). UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

Waste disposal: Search for and remove dead rodents at frequent intervals during treatment, at least as often as when baits are checked and/or replenished. Daily inspection may be required in some circumstances.

Dispose of dead rodents in accordance with local requirements.

In the UK poisoned rodents may be disposed of by the waste producer at an incinerator or landfill permitted to accept that type of waste, or collected by a registered waste carrier and taken for disposal at a suitably permitted site. For further information on disposal contact the Environment Agency (http://www.environment-agency.gov.uk/) or SEPA (http://www.sepa.org.uk/).

Remove all baits after treatment and dispose of them in accordance with local requirements. For further information on disposal contact the Environment Agency (http://www.environment-agency.gov.uk/) or SEPA (http://www.sepa.org.uk/).

Dead rodents and bait found outside bait stations should be disposed of using protective gloves.

Storage and shelf-life: Keep out of reach of children. Keep away from food and animal feedstuffs. Keep in a cool, dry, well-ventilated area. Store only in original container. Shelf life up to 2 years.

Authorisation Holder and Manufacturer:
Bábolna Bio Ltd., H-1107 Budapest, Szállás u. 6, Hungary
Tel: +36 1 432 0400 Fax: +36 1 432 0401
Email: info@babolna-bio.com , Internet: www.babolna-bio.com

® = *Registered trademark*

Manufacturing date/ Batch number:

S11751_2017_v01 UK

5997378771609